

Grand-Canonical simulation of 4D simplicial quantum gravity ^{*}

S.Horata^a, H.S.Egawa^b and T.Yukawa^c

^aDepartment of Particle and Nuclear Physics, School of Mathematical Science, The Graduate University for Advanced Studies, Tsukuba, Ibaraki 305-0801, Japan

^bDepartment of Physics, Tokai University, Hiratsuka, Kanagawa 259-1292, Japan

^cCoordination Center for Research and Education, The Graduate University for Advanced Studies, Hayama, Miura, Kanagawa 240-0193, Japan

A thorough numerical examination for the field theory of 4D quantum gravity (QG) with a special emphasis on the conformal mode dependence has been studied. More clearly than before, we obtain the string susceptibility exponent of the partition function by using the Grand-Canonical Monte-Carlo method. Taking thorough care of the update method, the simulation is made for 4D Euclidean simplicial manifold coupled to N_X scalar fields and N_A U(1) gauge fields. The numerical results suggest that 4D simplicial quantum gravity (SQG) can be reached to the continuum theory of 4D QG. We discuss the significant property of 4D SQG.

1. Introduction

Until now 4D SQG has been investigated from several points of view, such as the phase structure, nature of the transition, effects of additional matter fields, and the modified measure action [1]. Unfortunately the numerical results could not resolve the problem for the existence of the continuum theory of 4D QG clearly. Gradually the original enthusiasm to explore 4D QG has been toned down. However, we remind that many problems about 4D SQG are yet to be studied.

As one of those problems, we pay attention to additional matter effects in 4D Euclidean SQG (Eucl.SQG). According to the recent numerical simulations, the string susceptibility exponent ($\gamma^{(4)}$) takes a negative value with adding a few U(1) gauge fields. It makes drastic change of the phase diagram, and continuous phase transition with one gauge matter has been found[2]. Moreover, the numerically obtained $\gamma^{(4)}$ as a function of the number of matter fields turned out to be approximately equal to that of the prediction of 4D conformal field theory[3].

In order to see more definitely how the matter fields act on 4D Eucl.SQG, we repeat the study with employing the Grand-Canonical method

used for 2D SQG[4], which has given the satisfactory agreement of two results. For discussing the continuum theory of 4D QG based on the numerical analysis, we pay attention to the functional form of the partition function, which we expect enough to gain information for selecting the correct theory of gravity, if it exists at all. For obtaining the functional form of the partition function, we need to perform a simulation with varying the space volume. In practice, we calculate the probability ($P(N_4)$) as a function of the volume (N_4) by the Grand-Canonical method of the dynamical triangulation (DT) and estimate $\gamma^{(4)}$ from the scaling behavior of $P(N_4)$.

2. Model and Grand-Canonical method

Let us explain our model and the numerical method of the Grand-Canonical simulation. We consider a system of the simplicial gravity coupled to U(1) gauge fields and massless scalar fields. The action (S) is given on 4D simplicial manifold as

$$S = S_G + S_A + S_X, \quad (1)$$

where S_G , S_A and S_X denote the action for the gravity, U(1) gauge fields (A) and scalar fields (X), respectively.

^{*}presented by S.Horata

For the gravity part, we use the discretized Einstein-Hilbert action in 4D,

$$S_G[\kappa_2, \kappa_4] = \kappa_4 N_4 - \kappa_2 N_2, \quad (2)$$

where N_i denotes the number of i -simplex. The two parameters, κ_2 and κ_4 , correspond to the inverse of the gravitational constant and the cosmological constant, respectively. The action for the N_X scalar fields X_i^a , $a = 1 \dots N_X$, on the vertex i is given by

$$S_X = \sum_{a=1}^{N_X} \sum_{ij} o(l_{ij}) (X_i^a - X_j^a)^2, \quad (3)$$

where $o(l_{ij})$ is the number of four-simplices sharing a link l_{ij} . And the action for the N_A vector field A_{ij}^a ($a = 1 \dots N_A$) on link l_{ij} reads

$$S_A = \sum_{a=1}^{N_A} \sum_{ijk} o(t_{ijk}) (A^a(l_{ij}) + A^a(l_{jk}) + A^a(l_{ki}))^2, \quad (4)$$

where $o(t_{ijk})$ is the number of four-simplices sharing a triangle t_{ijk} . These actions give the correct thermo-dynamical limit for large N_4 because they are proportional to N_4 .

In order to simplify the calculation, the Monte-Carlo simulations have been performed with the Canonical method previously. This method requires that N_4 fluctuates around the given target volume (V_4). Thus, the volume fixing term (δS) is added to the total action Eq.(1),

$$\delta S(N_4, V_4) = \delta \kappa_4 (N_4 - V_4)^2, \quad (5)$$

where $\delta \kappa_4$ is the free parameter. However, the Canonical method has limitations about the finite size effect and the ergodic property. Furthermore, the functional form of the size dependence of the partition function cannot be given clearly. For the improvement of numerical accuracy we perform the simulations varying the space volume by the Grand-Canonical method. The Grand-Canonical partition function with the matter fields is given as

$$Z(\kappa_2, \kappa_4, N_A, N_X) = \sum_T e^{-S_G(\kappa_2, \kappa_4)}$$

$$\prod_{N_A, N_X} \left(\int \prod_{l \in T} dA_l e^{-S_A(A_l)} \int \prod_{i \in T} dX_i e^{-S_X(X_i)} \right), \quad (6)$$

where all geometries T and matter fields A, X are updated using the Metropolis method keeping the topology of discretized surface to be S^4 .

From the analogy of the 2D case, we assume that the partition function obeys the KPZ like formula for the large- N_4 limit as

$$Z \approx \sum_{N_4} N_4^{\gamma^{(4)}(\kappa_2) - 3} e^{(\kappa_4 - \kappa_4^c(\kappa_2)) N_4}. \quad (7)$$

Then the probability $P(N_4)$ for the large- N_4 behaves approximately as

$$P(N_4) \propto N_4^{\gamma^{(4)} - 3} e^{-(\kappa_4 - \kappa_4^c) N_4}. \quad (8)$$

By tuning κ_4 close to κ_4^c , the exponential growth is suppressed and $P(N_4)$ is given as the power function of N_4 . The parameter κ_4 is determined in the following manner: If κ_4 is chosen to be less (or greater) than κ_4^c , the system will expand (or shrink) exponentially in the Grand-Canonical simulation. By requiring an exponential increase (or decrease) to disappear in the simulation, we can find κ_4 precisely with the following form,

$$\frac{\ln P(N_4)}{\ln N_4} = (\gamma^{(4)} - 3) - (\kappa_4 - \kappa_4^c) \frac{N_4}{\ln N_4} + O(1/\ln N_4). \quad (9)$$

After tuning κ_4 , we generate the configurations constraining N_4 within an upper- and a lower-bound ($N_4^{min} \leq N_4 \leq N_4^{max}$).

3. Numerical result

In this section, we report numerical results with the Grand-Canonical method. In order to obtain $P(N_4)$, we allow N_4 to fluctuate between $N_4^{min} = 1000$ and $N_4^{max} = 10000$. In Fig.1, we show $P(N_4)$ versus N_4 at κ_4 close to κ_4^c . The numerical result shows that $P(N_4)$ is given as a power function of N_4 . It is suggested that the partition function in 4D obeys the KPZ type partition function as the 2D case.

We can also compute $\gamma^{(4)}$ from the slope of $P(N_4)$. Then, we show the parameter (b), which is calculated from γ^4 ,

$$\gamma^{(4)} = -\frac{b}{2} \left(1 - \frac{4}{b} + \sqrt{1 - \frac{4}{b}} \right). \quad (10)$$




Figure 1. We plot $P(N_4)$ versus N_4 with log-log scale.

This relation is given from the analogy of 2D QG and the parameter b is similar to the central charge in 2D QG. In Fig.2, we plot b versus the number of matter fields, $N_X + 62N_A$, on the critical point (κ_2^c) between the crumpled phase and the smooth phase. We compare b obtained by two different numerical methods, *i.e.* the Canonical method with the MINBU algorithm and the Grand-Canonical method. Error bars include both the errors in determining κ_2^c and the statistical errors, and we found that the former method has much larger errors than the latter method. We made the parameter fit of the data of the Grand-Canonical method by a linear function, $b = 0.0030(3)(N_X + 62N_A) + 3.98(3)$. The slope b is approximately equal to the analytical result[5,6].

4. Summary and Discussion

Let us summarize and discuss the main points made in the previous section. From the numerical results of the Grand-Canonical method, we find the KPZ type partition function also appears in 4D QG. The string susceptibility exponent $\gamma^{(4)}$ obtained from the size dependence of the partition function can be obtained by the Grand-Canonical method rather accurately. The matter contribution to b , which is computed from $\gamma^{(4)}$ with the analogy of 2D QG, turns out to be ap-




Figure 2. We plot b versus the number of the matter fields, $N_X + 62N_A$.

proximately equal to the conformal part[5,6]. Our numerical results suggest that it may be possible to understand the continuum theory of 4D QG from the analogy of 2D QG. In order to look more details on the relation to the continuum theory of 4D QG, these results encourage us to continue investigation by the Grand-Canonical method and push more efforts on the analytical calculation of the conformal gravity at the same time.

REFERENCES

- [1] G. Thorleifsson, Nucl. Phys. B (Proc. Suppl.) **73** (1998) 133.
- [2] H. S. Egawa, S. Horata, N. Tsuda and T. Yukawa, Prog. Theor. Phys. 106 (2001) 1037.
- [3] H. S. Egawa, S. Horata, and T. Yukawa, Nucl. Phys. B (Proc. Suppl.) **106-107** (2002), 971.
- [4] S. Oda, N. Tsuda and T. Yukawa, Nucl. Phys. B (Proc. Suppl.) **63** (1998) 733
- [5] H. S. Egawa, S. Horata, and T. Yukawa, Prog. Theor. Phys. submitted.
- [6] K. Hamada, Prog. Theor. Phys. 103 (2000) 1237, Prog. Theor. Phys. 105 (2001) 673.